

CPC GROWER R (BULK ONLY)

Medicated

230310

Active Drug Ingredient

Monensin (as Monensin sodium)29.4 g/ton

Guaranteed Analysis

Crude Protein	(Min)	13.00	%
Crude Fat	(Min)	2.00	%
Crude Fiber	(Max)	23.00	%

Ingredients

Processed Grain By-Products, Roughage Products, Grain Products, Calcium Carbonate and CPC PREMIX.

Feed to beef cattle on pasture. Feed at a rate of up to 1%-2% of body weight per head per day. Do not feed less than 3.4 pounds per head per day and not over 13.61 pounds per head per day to provide between 50 and 200 milligrams per head per day. Always feed with a free choice forage source. Always provide free choice minerals, vitamins and clean water. Do not offer free choice to any animal. Never feed wet or moldy feed. Always provide a source of fresh clean water, and free choice minerals. When making changes in an animal's diet, allow a 7 to 10-day transition period. Provide ample feeder space for livestock. Do not feed wet, moldy, or insect infested feed.

CAUTION

CAUTION: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulations containing monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result.

Limitations: For increased rate of gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed, or after the fifth day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of .14 to .42 milligrams per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first five days of feeding, cattle should receive not more than 100 milligrams per head in not less than 1 pound of feed.

This feed was developed for producers who wean and background their calf crop. We have seen excellent results when this grower feed is fed free choice to cattle from 400 to 800 pounds. The cattle develop at a healthy rate and develop their frame and muscling without adding too much fat and becoming over conditioned. Cattle grown out on CPC Grower have consistently been among the top selling cattle at several sales.

BULK ONLY