



PERF SHEEP PELL W/DECOX/AN

293020

Pelleted

Formulated to be fed to sheep

MEDICATED

For the prevention of coccidiosis in young sheep caused by *Eimeria bakuensis*, *Eimeria crandalis*, *Eimeria ovinoidalis* and *Eimeria parva*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT
Decoquinatate 14 g/ton

Guaranteed Analysis

Crude Protein	(Min)	18.00	%
Crude Fat	(Min)	3.00	%
Crude Fiber	(Max)	12.00	%
Calcium	(Min)	0.50	%
Calcium	(Max)	1.00	%
Phosphorus	(Min)	0.40	%
Vitamin A	(Min)	16,000	IU/lb
Vitamin D	(Min)	2,000	IU/lb
Vitamin E	(Min)	51	IU/lb

Ingredients

Processed Grain By-Products, Grain Products, Plant Protein Products, Forage Products, Roughage Products, Calcium Carbonate, Potassium Sulfate, Magnesium Sulfate, Ammonium Chloride, Kaolin, Salt, Expand, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement and Sacchanomyces Yeast Culture.

Feeding Directions

This feed is formulated for sheep and should be fed at a rate of 2% of body weight per head per day.

CAUTION

WARNING: Do not feed to sheep producing milk for food.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed Type C feed or milk replacer at a rate to provide 22.7 mg per 100 lb (0.5 mg/kg) bodyweight. Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid type B feed.

Always provide a source of fresh water, and free choice minerals. When making changes in an animals diet, allow a 7 to 10 day transition period. Provide ample feeder space for livestock. Do not feed wet, moldy, or insect infested feed

MEDICATED

This is a great feed for lambs and goat kids, both as a creep feed and for the growth and finishing phases. It is high energy and high protein and good weight gains can be expected.

It is also a good option for the show lambs and goats.